

Health Watch Table — Down Syndrome

Forster-Gibson and Berg 2011

CONSIDERATIONS	RECOMMENDATIONS
1. HEENT (HEAD, EYES, EARS, NOSE, THROAT)	
<p>Children and Adults: Vision: ~15% have cataracts; ~ 20% - 70% have significant refractive errors</p> <p>5% - 15% of adults have keratoconus</p> <p>Hearing: 50% - 80% have a hearing deficit</p>	<ul style="list-style-type: none"> <input type="checkbox"/> Neonatally: refer immediately to an ophthalmologist if the red reflex is absent or if strabismus, nystagmus or poor vision is identified. <input type="checkbox"/> Arrange ophthalmological assessment: first by 6 months for all; then every 1-2 years, with special attention to cataracts, keratoconus, and refractive errors. <input type="checkbox"/> During childhood: screen vision annually with history and exam; refer as needed. <input type="checkbox"/> Arrange auditory brainstem response (ABR) measurement by 3 months if newborn screening has not been done or if results were suspicious. <input type="checkbox"/> During childhood: screen hearing annually with history and exam; review risks for frequently occurring serious otitis media. <input type="checkbox"/> Undertake auditory testing: first at 9 – 12 months, then every 6 months up to 3 years, annually until adulthood, then every two years.
2. DENTAL	
<p>Children and Adults: Tooth anomalies are common</p> <p>Increased risk of periodontal disease in adults</p>	<ul style="list-style-type: none"> <input type="checkbox"/> Undertake initial dental exam at 2 years, then every 6 months thereafter. Encourage proper dental hygiene. Refer to an orthodontist if needed. <input type="checkbox"/> Undertake clinical exams every six months with referral, as appropriate.
3. CARDIOVASCULAR	
<p>Children: 30% - 60% have congenital heart defects (CHD)</p>	<ul style="list-style-type: none"> <input type="checkbox"/> Newborn screening: Obtain an echocardiogram and refer to a cardiologist, even in the absence of physical findings. <input type="checkbox"/> In children and adolescents: review cardiovascular history and assess for physical signs with specialist referral, if indicated. <ul style="list-style-type: none"> • Refer for an echocardiogram if not previously done • Undertake SBE prophylaxis as indicated by findings
<p>Adults: ~ 50% have cardiovascular concerns, commonly acquired mitral valve prolapse (MVP) and valvular regurgitation</p>	<ul style="list-style-type: none"> <input type="checkbox"/> Ascertain a comprehensive cardiovascular history. <input type="checkbox"/> Undertake an annual cardiac exam, with echocardiogram to confirm new abnormal findings and follow-up depending on the type of cardiovascular problem present or refer to an Adult Congenital Heart specialist or Disease clinic. <input type="checkbox"/> Monitor regularly those that have had surgery in childhood. <input type="checkbox"/> An echocardiogram is indicated to assess new abnormal physical findings or if unable to assess adequately by physical exam. Consider echocardiogram to establish baseline cardiac anatomy and function if not previously done or records are unavailable.¹
4. RESPIRATORY	
<p>Children and Adults: 50% - 80% have obstructive sleep apnea (OSA)</p>	<ul style="list-style-type: none"> <input type="checkbox"/> Newborn: Refer to an ENT surgeon if recurring otitis media infections. <input type="checkbox"/> Treat infections promptly and aggressively.
<p>Adults: 50% - 80% have obstructive sleep apnea (OSA)</p>	<ul style="list-style-type: none"> <input type="checkbox"/> Ascertain a detailed sleep history, with special attention to OSA symptoms. Refer to an ENT surgeon, including sleep study, if OSA is suspected. <input type="checkbox"/> If aspiration pneumonia is suspected, investigate for possible swallowing disorder and gastro-esophageal reflux disease.

CONSIDERATIONS	RECOMMENDATIONS
5. GASTROINTESTINAL	
Children: ~ 50% have gastrointestinal (GI) tract anomalies including duodenal atresia, celiac disease, Hirschsprung disease, and imperforate anus	<ul style="list-style-type: none"> <input type="checkbox"/> Newborn: with vomiting or absent stools, check for GI tract blockage and refer to a gastroenterologist. <input type="checkbox"/> Infants and children: anticipate constipation; treat with fluid/fibre/laxative/stool softener/exercise/dietary change. <input type="checkbox"/> From 2-3 years of age, screen for celiac disease. <input type="checkbox"/> Establish good dietary and exercise habits to prevent or manage obesity.
Adults: ~ 95% are obese; ~ 7% have celiac disease	<ul style="list-style-type: none"> <input type="checkbox"/> Monitor for obesity. <input type="checkbox"/> Screen for celiac disease, which may present in adulthood; screening tests used are the same as in the general population.² <input type="checkbox"/> Test for Helicobacter Pylori and treat if positive, regardless of symptoms. <input type="checkbox"/> Manage constipation proactively.
6. GENITOURINARY	
Children: Cryptorchidism is common	<ul style="list-style-type: none"> <input type="checkbox"/> Assess for hypogonadism, undescended testes, and possible testicular germ-cell tumors, or refer to a urologist, as appropriate.
Adults: Have increased risk of testicular cancer	<ul style="list-style-type: none"> <input type="checkbox"/> Assess annually by clinical exam, and refer to a urologist as appropriate.³
7. SEXUAL FUNCTION	
Adults: Fertility has been documented in women Fertility in males rarely reported	<ul style="list-style-type: none"> <input type="checkbox"/> Counsel regarding fertility possibility and the 50%⁴ risk of Down syndrome in offspring.
8. MUSCULOSKELETAL (MSK)	
Children: ~15% have atlantoaxial instability (AAI)	<ul style="list-style-type: none"> <input type="checkbox"/> Arrange lateral cervical spine X-rays (flexed, neutral, and extended positions) between 3-5 years of age. <input type="checkbox"/> Screen, as needed, prior to high risk activities (e.g., tumbling) and if participating in Special Olympics. <input type="checkbox"/> Undertake an annual neurological exam for signs or symptoms of spinal cord compression. If present, refer urgently to a neurosurgeon and arrange an urgent MRI. <input type="checkbox"/> Obtain a detailed MSK history with particular attention to possible joint subluxations/dislocations, scoliosis, and hip abnormalities.
Adults: Continued risk for spinal cord compression secondary to AAI Though data are limited, osteoporosis (associated with increased fractures risk) may be more common in older adults with Down syndrome than in similar aged individuals in the general population or with other developmental disabilities	<ul style="list-style-type: none"> <input type="checkbox"/> Undertake an annual neurological exam and assess for evidence of spinal cord compression. <input type="checkbox"/> Arrange lateral cervical spine X-rays if not previously done, if presenting with signs and symptoms of AAI or if participating in Special Olympics. <input type="checkbox"/> Take detailed history and attend to joint complaints, scoliosis, and hip abnormalities. <input type="checkbox"/> If suspected, undertake bone mineral density (BMD) screening and refer to an appropriate specialist, if indicated. <input type="checkbox"/> Encourage ambulation/mobility and weight reduction if obesity is present to decrease the risk of osteoarthritis.
9. NEUROLOGY	
Children: Epilepsy in up to 22%	<ul style="list-style-type: none"> <input type="checkbox"/> Take careful neurological history with particular attention to seizures (infantile spasms or tonic-clonic-type). <input type="checkbox"/> Arrange an EEG and refer to a neurologist.

CONSIDERATIONS	RECOMMENDATIONS
<p>Adults: Dementia is frequent and occurs earlier: 11%: 40 – 49 y, 77%: 60 – 69 y, Up to 75% with dementia have seizures with frequency increasing with age</p>	<ul style="list-style-type: none"> <input type="checkbox"/> Obtain a neuropsychiatric history at every visit with particular attention to change in behaviour, loss of function/activities of daily living, and new onset seizures. <input type="checkbox"/> If functional decline and/or signs/symptoms of dementia, use history, exam, and blood work to check for other conditions and treatable causes (e.g., hearing/vision deficits, obstructive sleep apnea, hypothyroidism, chronic pain, medication side effects, depression, menopause, low folic acid/vitamin B12). <input type="checkbox"/> For possible seizures, arrange an EEG and refer to a neurologist.
10. DERMATOLOGICAL	
<p>Children and Adults: Dry skin, atopic dermatitis, seborrheic dermatitis, chelitis, impetigo, and alopecia areata are more common than in general population</p>	<ul style="list-style-type: none"> <input type="checkbox"/> Examine skin as part of routine care. <input type="checkbox"/> Treat as per general population, with referral to dermatologist as needed.
11. BEHAVIOIURAL/MENTAL HEALTH	
<p>Children: Self-talk is very common; autism spectrum disorder occurs in 5% - 10% of children with DS</p>	<ul style="list-style-type: none"> <input type="checkbox"/> Review regularly with respect to behavioural concerns. <input type="checkbox"/> Review for positive or negative signs suggestive of psychosis.
<p>Adults: ~ 30% have a psychiatric disorder, including depression</p>	<ul style="list-style-type: none"> <input type="checkbox"/> Review regularly with respect to behavioural concerns. <input type="checkbox"/> Ascertain neuropsychiatric history at every visit, with particular attention to changes in behaviour, loss of function/activities of daily living, and new onset seizures.
12. ENDOCRINE	
<p>Children: ~ 1% have congenital hypothyroidism; ~ 20% develop hypothyroidism after birth</p>	<ul style="list-style-type: none"> <input type="checkbox"/> Review neonatal screening. <input type="checkbox"/> Ascertain TSH and free T4 tests to confirm euthyroid status at 6 and 12 months, then annually. <input type="checkbox"/> If signs of hyperthyroidism in adolescence, check for autoimmune thyroiditis.
<p>Adults: 15% - 50% are hypothyroid Subclinical hypothyroidism, hyperthyroidism, and autoimmune thyroiditis are more common than in the general population</p>	<ul style="list-style-type: none"> <input type="checkbox"/> For adults who are euthyroid, check TSH and free T4 levels at least once every 5 years ⁵ (some recommend annually).⁶ <input type="checkbox"/> If subclinical hypothyroidism (i.e., elevated TSH with normal free T4), follow free T4 every 6 months for one year ⁷ (some recommend treatment if thyroid antibodies are positive). <input type="checkbox"/> Consider checking thyroid function whenever there are changes in mental status, behaviour or functional abilities.
13. HEMATOLOGICAL	
<p>Children and Adults: Increased frequency of transient myeloproliferative disorder and leukemia No increased risk of leukemia in adults</p>	<ul style="list-style-type: none"> <input type="checkbox"/> Neonates to 1 month olds: investigate for polycythemia and thromobocytopenia. <input type="checkbox"/> Assess history periodically for symptoms of leukemia, with close attention to those with a history of transient myeloproliferative disorder.

WEBSITES THAT MAY BE HELPFUL FOR FAMILIES AND CAREGIVERS

Canadian Down Syndrome Society	<input type="checkbox"/> www.cdss.ca/
Down Syndrome Education International [DownsEd]	<input type="checkbox"/> www.downsed.org/
Down Syndrome: Health Issues by Dr. Len Leshin	<input type="checkbox"/> www.ds-health.com/
Down Syndrome Medical Interest Group [DSMIG-UK]	<input type="checkbox"/> www.dsmig.org.uk/
National Down Syndrome Society [USA]	<input type="checkbox"/> www.ndss.org/

REFERENCES CITED

1. Lin AE, Basson CT, Goldmuntz E, Magoulas PL, McDermott DA, McDonald-McGinn DM, et al. Adults with genetic syndromes and cardiovascular abnormalities: clinical history and management. *Genet Med* 2008 Jul;10(7):469-94.
2. National Institute for Clinical Excellence [NICE]. Coeliac disease: recognition and assessment of coeliac disease – Quick reference guide. Nice clinical guideline 86. London: National Institute for Clinical Excellence [NICE]; 2009. Available from www.nice.org.uk/nicemedia/pdf/CG86QuickRefGuide.pdf.
3. Patja K, Pukkala E, Sund R, Iivanainen M, Kaski M. Cancer incidence of persons with Down syndrome in Finland: a population-based study. *Int J Cancer* 2006 Apr 1;118(7):1769-72.
4. Galley R. Medical management of the adult patient with Down syndrome. *JAAPA*. 2005 Apr;18(4):45,6, 48, 51-2.
5. Prasher V, Gomez G. Natural history of thyroid function in adults with Down syndrome--10-year follow-up study. *J Intellect Disabil Res* 2007 Apr;51(Pt 4):312-7.
6. McGuire D, Chicoine B. Chapter 2: Assessing the physical health/mental health connection. In: McGuire D, Chicoine B, editors. *Mental wellness in adults with Down syndrome: A guide to emotional and behavioral strengths and challenges*. Bethesda, MD: Woodbine House; 2006. p. 9-28.
7. Wallace RA, Dalton AJ. Clinicians' guide to physical health problems of older adults with Down syndrome. *Journal on Developmental Disabilities* 2006;12 (1 [Supplement 1]):1-92. Available from www.oadd.org/publications/journal/issues/vol12no1supp/DS_supplement_1.pdf.

PUBLISHED HEALTH CARE GUIDELINES REVIEWED AND COMPARED

- American Academy of Pediatrics. Committee on Genetics, American Academy of Pediatrics: Health supervision for children with Down syndrome. *Pediatrics*, 2001, 107, 2, 442-449, United States.
- Baum, R.A.; Nash, P.L.; Foster, J.E.; Spader, M.; Ratliff-Schaub, K.; Coury, D.L., Primary care of children and adolescents with down syndrome: an update. *Curr.Probl.Pediatr.Adolesc.Health.Care.*, 2008, 38, 8, 241-261, United States.
- Bosch, J.J., Health maintenance throughout the life span for individuals with Down syndrome. *J.Am.Acad.Nurse Pract.*, 2003, 15, 1, 5-17, United States.
- Cohen, W.I.; The Down Syndrome Medical Interest Group (DSMIG), Health care guidelines for individuals with Down syndrome: 1999 Revision (Down syndrome preventive medical checklist). *Down Syndrome Quarterly*, 1999, 4, 3, 1-16.
- Cooley, W.C.; Graham, J.M., Jr., Down syndrome – an update and review for the primary pediatrician. *Clin.Pediatr.(Phila)*, 1991, 30, 4, 233-253, United States.
- Davidson, M.A., Primary care for children and adolescents with Down syndrome. *Pediatr.Clin.North Am.*, 2008, 55, 5, 1099-1111, United States.
- Galley, R., Medical management of the adult patient with Down syndrome. *JAAPA*, 2005, 18, 4, 45-6, 48, 51-2, United States.
- Henderson, A.; Lynch, S.A.; Wilkinson, S.; Hunter, M., Adults with Down's syndrome: the prevalence of complications and health care in the community. *Br.J.Gen.Pract.*, 2007, 57, 534, 50-55, England, Guideline 00 - Introduction & Discussion.
- Hunter, Alasdair, *Down syndrome*. 2005, 191-210, Wiley-Liss, Hoboken, N.J., 2nd.
- Pueschel, Siegfried M., *Optimal health care and medical concerns*. 2006, 9-46, Paul H. Brookes Pub. Co., Baltimore, Md.

- Saenz, R.B., Primary care of infants and young children with Down syndrome. *Am.Fam.Physician*, 1999, 59, 2, 381-90, 392, 395-6, United States.
- Smith, D.S., Health care management of adults with Down syndrome. *Am.Fam.Physician*, 2001, 64, 6, 1031-1038, United States.
- van Allen, M.I.; Fung, J.; Jurenka, S.B., Health care concerns and guidelines for adults with Down syndrome. *Am.J.Med.Genet.*, 1999, 89, 2, 100-110, Wiley-Liss, Inc, United States.
- Van Cleve, S.N.; Cohen, W.I., Part I: Clinical practice guidelines for children with Down syndrome from birth to 12 years. *J. Pediatr. Health Care*, 2006, 20, 1, 47-54.
- Van Cleve, S.N.; Cannon, S.; Cohen, W.I., Part II: Clinical Practice Guidelines for Adolescents and Young Adults With Down Syndrome: 12 to 21 Years. *J. Pediatr. Health Care*, 2006, 20, 3, 198-205.
- Wallace, Robyn A.; Dalton, Arthur J., Clinicians' guide to physical health problems of older adults with Down syndrome. *Journal on Developmental Disabilities*, 2006, 12, 1 (Supplement 1), 1-92.
- Wilson, Golder N.; Cooley, Carl W., Preventive management of Down syndrome. 2006, 175-193, Cambridge University Press, Cambridge, 2nd.

Developed by: **Forster-Gibson, Cynthia, MD, PhD and Berg, Joseph M, MB, BCh, MSc, FRCPSYCH, FCCMG**

Expert Clinician Reviewers

Thanks to the following clinicians for their review and helpful suggestions.

Brian Chicoine, MD
 Medical Director, Adult Down Syndrome Center
 of Lutheran General Hospital
 Park Ridge, Illinois

Len Leshin, MD
 Down Syndrome: Health Issues
 Corpus Christi, Texas

About this Health Watch Table

Initial publication:
 May 2011

Edit history:
 20 November 2012, PT
 26 May 2015, PT