



GLOBAL
DOWN SYNDROME FOUNDATION®



GLOBAL DOWN SYNDROME FOUNDATION
MEDICAL CARE GUIDELINES
for ADULTS WITH DOWN SYNDROME

Family-Friendly Version

Introduction

1 Behavior

2 Dementia

3 Diabetes

4 Cardiovascular Disease

5 Obesity

6 Atlantoaxial Instability

7 Osteoporosis

8 Thyroid

9 Celiac Disease

Appendix & References

AUTHORS

Peter Bulova, MD

George Capone, MD

Brian Chicoine, MD

Bryn Gelaro, MA, LSW

**Terry Odell Harville, MD, PhD,
D(ABMLI), F(ACHI)**

Barry A. Martin, MD

**Dennis E. McGuire,
PhD, LCSW**

***Kent D. McKelvey, MD**

**Moya Peterson, PhD,
APRN, FNP-BC**

Amy Tsou, MD, MSc

**Carl Tyler, MD, MSc, ABFP,
CAQ-Geriatrics**

Michael David Wells, BS

Michelle Sie Whitten, MA





VOLUNTEER COMMITTEE

Jarrett Barnhill, MD, DFAPA, FAACAP
Donald Bodenner, MD, PhD
Paul J. Camarata, MD
Kamala Gullapalli Cotts, MD
Robert H. Eckel, MD
Anna J. Esbensen, PhD
James E. Hunt, MD
Seth M. Keller, MD
Judy Lu Kim, MD
Ira T. Lott, MD
Michael T. McDermott, MD
Joan Medlen, MEd, RD, LD
Micol Rothman, MD
Stephanie L. Santoro, MD
Mary M. Stephens, MD, MPH
Elizabeth Yeung, MD

OTHER EXPERTS

Joaquín M. Espinosa, PhD
Joann Fontanarosa, PhD
Gina Giradi, MS
Allison Gross, MLS
Janice Kaczmarek, MS
Lina Patel, PsyD
Michael S. Rafii, MD, PhD
Karen Schoelles, MD, SM, FACP

KEY COLLABORATORS

The Ritter Family

BENEFACTORS

Burg Family Foundation

Down Syndrome Association
of Greater Richmond

Down Syndrome Guild of
Dallas

National Down Syndrome
Society

Rocky Mountain
Down Syndrome Association

MULTI-YEAR SUPPORTERS

arc Thrift Store

Down Syndrome Alliance
of the Midlands

Down Syndrome Association
of Atlanta

Down Syndrome Association
of Greater St. Louis

Down Syndrome Association
of Middle Tennessee

Down Syndrome Association
of Minnesota

Down Syndrome Association of
Northeast Ohio

Down Syndrome Association of
the Bay Area

Dra. Macarena Lizama & Centro US
Síndrome de Down

Japan Down Syndrome Association

North Carolina Down Syndrome
Alliance

KEY SUPPORTERS

Anna & John J. Sie Center
for Down Syndrome

Charles Monfort

Down Syndrome Association
of Central Ohio

Down Syndrome Association
of Jacksonville


Down Syndrome Association
of Orange County

Down Syndrome of Louisville, Inc

Linda Crnic Institute for
Down Syndrome

National Down Syndrome Congress

The Napoleone Family



SUPPORTERS

Broward Gold Coast
Down Syndrome Organization

ChapTer 21

Designer Genes of North Dakota

Down Syndrome Affiliates in Action

Down Syndrome Alabama

Down Syndrome Association for
Families of Nebraska

Down Syndrome Association of
Acadiana

Down Syndrome Association of
Central Florida

Down Syndrome Association of
Connecticut

Down Syndrome Association of
Greater Cincinnati

Down Syndrome Association of
Northern Virginia

Fun Coast Down Syndrome
Association

The Family of Rya Gracyn Pierce

FRIENDS

Alan P. & Veronica C. Neuman
Family Charitable Fund

Alaska Down Syndrome Network

Chattanooga Down Syndrome
Society

Club 21 Learning and Resource
Center

David Egan and Family

Down Syndrome Association
of Central California

Down Syndrome Association
of Central Kentucky

Down Syndrome Association
of Central Oklahoma

Down Syndrome Association
of Central Texas

Down Syndrome Association
of Delaware

Down Syndrome Association
of the Brazos Valley

Down Syndrome Association of
Wisconsin, Inc.

Down Syndrome Family
Connection

Down Syndrome Foundation
of Southeastern New Mexico

Eastern Idaho Down Syndrome

Family Connect East Texas Down
Syndrome Group

Families Exploring
Down Syndrome of Brevard

International Mosaic
Down Syndrome Association

Kern Down Syndrome Network

Red River Valley
Down Syndrome Society

Rio Grande Valley
Down Syndrome Association

Sampson Collaborative Law

S.M.I.L.E. on Down Syndrome

Southern Arizona Network
for Down Syndrome

Wisconsin Upside Down



INTRODUCTION


People with Down syndrome have different medical needs than people without Down syndrome.

For example, there are some medical conditions they are more likely to have, and some conditions they are less likely to have, compared to people without Down syndrome. So, their annual check-ups and sometimes their course of treatments may be different.

There are healthcare guidelines for children with Down syndrome ages 0-21 that are updated from time to time,^{1,2} which are very different from healthcare guidelines for children without Down syndrome. Although healthcare guidelines for adults with Down syndrome were written in 1999³ and 2001,⁴ they were based only on clinician experience and were not updated, even though there have since been advances and changes to standards of care.

In 2014, the Global Down Syndrome Foundation (GLOBAL) organized a task group of adults with Down syndrome, their families/caregivers, and Down syndrome experts for the purpose of establishing a medical care center for adults with Down syndrome. The feedback from that task group identified medical guidelines for adults with Down syndrome as an important tool for medical clinics and healthcare professionals to help provide more accurate evaluation and care.

As a result, the Global Down Syndrome Foundation Medical Care Guidelines for Adults with Down Syndrome Workgroup (GLOBAL Workgroup) was formed in 2016 to create these first ever evidence-based guidelines for adults with Down syndrome to improve the health of adults with Down syndrome. The Global Medical Care Guidelines for Adults with Down Syndrome (GLOBAL Adult Guideline) was published in JAMA (Journal of the American Medical Association) in October 2020.⁵ That was a very important milestone since medical professionals rely on trusted journals like JAMA and can easily find and implement the guidelines for their adult patients with Down syndrome. The GLOBAL Adult Guideline published in JAMA (2020) can be found online here: <https://bit.ly/GLOBAL-Adult-Guideline-JAMA>



This is the family-friendly version of the GLOBAL Adult Guideline. The original, full length GLOBAL Adult Guideline was written for a highly technical, medical audience. A version specifically for self-advocates is a future project. The recommendations in the family-friendly version are the same, but the content has been modified, with research components greatly abbreviated, and the technical methodology mostly removed. The original GLOBAL Adult Guideline can be found online here: <https://www.globaldownsyndrome.org/medical-care-guidelines-for-adults/>.

All images are approved GLOBAL models.



WHAT IS DOWN SYNDROME?

There Are Three Types Of Down Syndrome:

TRISOMY 21 About 95% of people with Down syndrome have Trisomy 21⁶ – a condition in which a person is born with three copies of chromosome 21 instead of the usual two copies in every cell.

TRANSLOCATION DOWN SYNDROME About 3% of people with Down syndrome have Translocation Down syndrome. This is when part or all of chromosome 21 attaches (translocates) onto another chromosome.

MOSAIC DOWN SYNDROME About 2% of people with Down syndrome have Mosaic Down syndrome.⁶ This is when only some (not all) of a person's cells have an extra copy of chromosome 21.

Down syndrome is the most common chromosomal condition⁷ and cause of developmental delay in the United States. Down syndrome is a genetic condition, not a disease. About 1 in every 691 babies born has it.⁸ Most of the time, Down syndrome is random and not inherited.⁹ How and why people have Down syndrome is largely unknown.

SINCE IT AFFECTS ALL OF THE BODY'S SYSTEMS, PEOPLE WHO HAVE DOWN SYNDROME ARE MORE LIKELY TO GET CERTAIN DISEASES THAN PEOPLE WITHOUT DOWN SYNDROME, AND LESS LIKELY TO GET OTHERS, SO THEY NEED SPECIAL MEDICAL CARE.

People with Down syndrome are less likely to have solid malignancies (cancerous tumors) or atherosclerotic cardiovascular disease (clogged arteries) than people without Down syndrome.^{10,11} The medical problems that people with Down syndrome are more likely to have include autoimmune conditions, leukemia, respiratory infections, sleep apnea, hearing and vision loss, and dementia. This does not mean that everyone with Down syndrome will have any or all of these problems. In addition, most of these problems can be found and treated early, which means people with Down syndrome who have them can still enjoy a good quality of life.¹²

Most people with Down syndrome have some intellectual disability.² Down syndrome affects everyone differently, but some common features include limited language skills, delayed muscle coordination, and reduced executive functioning. Other social and environmental difficulties, like finding knowledgeable healthcare or lack of access, may make things even harder and should also be taken into account as part of the care for adults with Down syndrome.

ABOUT THIS GUIDELINE

The GLOBAL Adult Guideline focuses on adults over age 21 with the type of Down syndrome called Trisomy 21. Since Down syndrome affects every part of the body,¹⁷ it was not possible to cover every medical area in this first edition of the guidelines. The GLOBAL Workgroup decided to focus on nine of the most common and concerning issues, which are:

- | | | |
|-------------|-----------------------------|-------------------|
| 1. Behavior | 4. Cardiovascular Disease | 7. Osteoporosis |
| 2. Dementia | 5. Obesity | 8. Thyroid |
| 3. Diabetes | 6. Atlantoaxial Instability | 9. Celiac Disease |

Since there are still many other medical areas that we believe are important and unique to people with Down syndrome, GLOBAL has committed to invest the time, energy and funds to update the GLOBAL Adult Guideline every 5-6 years with new medical areas and updates to the original nine. Currently, the GLOBAL Workgroup is also researching sleep apnea, solid tumors, leukemia, other autoimmune disorders, vision and eye issues, and physical therapy and fitness.

As a direct result of GLOBAL's advocacy in Washington, DC, the amount of funding for Down syndrome research increased from \$27 million in 2016 to \$115 million in 2022. GLOBAL expects this increase in funding will continue, and we will keep asking for support to elongate life and improve health outcomes with an additional focus on adults with Down syndrome. In the future, we also want to study the dual diagnosis of Autism and Down syndrome, Mosaic Down syndrome, and health disparities across gender and race in adults with Down syndrome.

PREVALENCE & LIFE COURSE

In the US, people with Down syndrome are living longer now than ever before. The lifespan of a person with Down syndrome has increased from 25 years in the 1980's¹³ to 60 years today.¹⁴

People with Down syndrome are living longer because they are no longer institutionalized, and there have been great improvements in treating pediatric heart problems, hypothyroidism and lung disease, as well as better access to medical care, education, community support, and other opportunities. There are also more babies being born with Down syndrome.^{8,15}

The government doesn't keep track of how many people have Down syndrome, so nobody knows the total population for sure. An estimate from 2010 said that there are 206,000 people with Down syndrome in the United States, which includes 125,000 adults.¹⁶ If we update those figures for today, there could be over 400,000 people with Down syndrome.

Even though there are more people with Down syndrome living longer, we need to do a lot more work in the areas of research and medical care to help keep people with Down syndrome healthy throughout adulthood.

GLOBAL ADULT GUIDELINE CHALLENGES

The biggest challenge when writing the GLOBAL Adult Guideline was not finding enough good research on Down syndrome or research that includes people with Down syndrome. More research is needed to support the way medical care is actually provided and in order to make better guideline recommendations. This research needs to be funded by the National Institutes of Health (NIH) along with universities, hospitals and any other advocacy and medical organizations in the Down syndrome community. Even without that research, the GLOBAL Workgroup was able to provide sound recommendations based on decades of clinical experience and the limited research available. When there wasn't enough research on adults with Down syndrome, we looked at research from other groups of people, such as children with Down syndrome or adults with other disabilities to decide if those guidelines could apply.¹⁸

METHODS

What makes the GLOBAL Adult Guideline first-in-kind is that it is evidence-based, meaning the authors reviewed research articles, identified articles they determined were of high quality, and then based their recommendations off what they found in the research. To be evidence-based, the GLOBAL Workgroup recruited guideline experts at ECRI and followed a very specific best practice called the GRADE Evidence to Decision Making Framework.

Once the guideline was written, we asked adults with Down syndrome and their family members for feedback via an online survey. GLOBAL also asked medical professionals from the American Academy of Developmental Medicine and Dentistry to review the draft guideline and provide feedback. Based on the survey and the feedback, the GLOBAL Workgroup made some important changes to the GLOBAL Adult Guideline before it was published.

To review the complete methodology, please visit the full GLOBAL Adult Guideline here:
<https://www.globaldownsyndrome.org/medical-care-guidelines-for-adults/>



1

BEHAVIOR



★ Statement of Good Practice 1 A review of behavioral, functional, adaptive, and psychosocial factors should be performed as part of an annual history that clinicians obtain from all adults with Down syndrome, their families, and caregivers.

Adults with Down syndrome should get the same level of medical care as adults without Down syndrome. This should include an annual review of all relevant behavioral (mood, attention, and activity level), functional-adaptive (social, communication, and daily living skills), and psychosocial (major life events, loss/grief, and trauma) factors that may affect physical and mental well-being. Since adults with Down syndrome may not be able to explain these things well, the family, caregivers and significant others should tell medical professionals about anything they've noticed that has changed in the mood or behavior of the adult with Down syndrome. If there are no big changes, that information can be used to establish a baseline.

★ Statement of Good Practice 2 When concern for a mental health disorder in adults with Down syndrome is present, medical professionals should evaluate for medical conditions that may present with psychiatric and behavioral symptoms.

Behavior and mental health conditions are common in adults with Down syndrome¹⁹ and are a cause for concern with families and caregivers.

WHEN THERE IS A CHANGE IN THE ADULT WITH DOWN SYNDROME'S BEHAVIOR OR FUNCTION, IT IS BEST TO HAVE A MEDICAL AND MENTAL HEALTH EVALUATION THAT LOOKS AT OTHER MEDICAL PROBLEMS THAT COULD CAUSE THOSE SYMPTOMS, SINCE SOMETIMES MEDICAL CONDITIONS MAY LOOK LIKE A BEHAVIORAL ISSUE.²⁰⁻²³


For example, head-banging when a person has a migraine, or not participating in an activity because they feel sick in their stomach.

RECOMMENDATION 1

When concern for a mental health disorder in adults with Down syndrome is present, medical professionals should refer to a clinician knowledgeable about the medical, mental health disorders, and common behavioral characteristics of adults with Down syndrome.

Behavior and mental health conditions are common in adults with Down syndrome¹⁹ and are a cause for concern with families and caregivers. An accurate diagnosis of a mental health disorder is important so as to provide quick and effective treatment. It is also important as to avoid misdiagnosing adaptive behavior as a disorder and providing the incorrect treatment. Assessment for underlying medical conditions significantly improves understanding the causes for changes in behavior or mental health and improves treatment.

We did not find any appropriate evidence-based studies that compare symptoms of mental health conditions in adults with Down syndrome to adults without Down syndrome. However, medical professionals who do not have a lot of experience in caring for adults with Down syndrome may find it challenging to treat them correctly. For example, adults with Down syndrome are very sensitive to certain drugs. Treatment is better if medical professionals know about the drug tolerances and intolerances people with Down syndrome have and how to balance those differences. It can be hard to find a medical professional who knows about these differences, but you can find the Down syndrome clinic nearest you here: <https://www.globaldownsyndrome.org/research-medical-care/medical-care-providers/>




RECOMMENDATION 2

When concern for a mental health disorder in adults with Down syndrome is present, medical professionals should follow guidelines for diagnosis in the Diagnostic and Statistical Manual of Mental Disorders (5th Ed; DSM-5).²⁴ The Diagnostic Manual-Intellectual Disability 2 (DM-ID-2)²⁵ also may be used to adapt diagnostic criteria from the DSM-5.

It can be challenging to diagnose mental health conditions in adults with Down syndrome because their behavior may look different from adults without Down syndrome. These differences may include limited speaking skills, difficulty adapting to changes, having more than one medical problem, and not being able to see a medical professional who is familiar with mental health in adults with Down syndrome. In our experience, adults with Down syndrome frequently use self-talk, have vivid imaginary/fantasy worlds, and stick to certain routines. Certain types of behavior may be common in adults with Down syndrome (e.g., self-talk), while other behaviors may be a sign of a treatable mental health condition (e.g., agitation, irritability). Medical professionals should be careful about assuming these behaviors are part of a mental health disorder in adults with Down syndrome. Instead of relying on an adult with Down syndrome to self-report their emotional problems, it is important for medical professionals to know that maladaptive behavior may be an indicator.

If skills decline in an adult with Down syndrome, it should be looked into. Loss of function may be a sign of a medical or mental health disorder. Diagnosis should take into consideration common behaviors and be based on observed behavior in addition to self-reporting.²⁶⁻³⁰



2

DEMENTIA





RECOMMENDATION 3

Caution is needed when diagnosing age-related, Alzheimer's-type dementia in adults with Down syndrome younger than age 40 due to its low prevalence before this age.

Dementia describes the symptoms of memory loss and changes in language, thinking, and/or functional skills. Alzheimer's disease is a type of a dementia. This recommendation is intended to decrease the misdiagnosis of age-related, Alzheimer's-type dementia in adults with Down syndrome. Misdiagnosis of age-related dementia in adults with Down syndrome may occur if medical professionals do not carefully evaluate for other medical conditions that can also cause functional decline. Adults with Down syndrome are more likely to have dementia, but it is uncommon to show signs before age 40. It is important for medical professionals working with patients of all ages to consider other causes to avoid misdiagnosis of Alzheimer's-type dementia. Misdiagnosis keeps people from getting correct treatment. We support careful and thorough medical evaluation to see if there are other causes of dementia-type symptoms in adults with Down syndrome under 40 years old. While there are no medical treatments for dementia, there are many guidelines or treatments for helping with behaviors, quality of life, and depression, which may occur.



RECOMMENDATION 4

Medical professionals should assess adults with Down syndrome and interview their primary caregivers about changes from baseline function annually beginning at age 40. Decline in the following six domains, as per the National Task Group – Early Detection Screen for Dementia (NTG-EDSD),³¹ should be used to identify early-stage age-related Alzheimer’s-type dementia and/or a potentially reversible medical condition:

- Cognition, memory, and executive function
- Behavior and personality
- Communication
- Adaptive functioning
- Ambulation and motor skills
- General decline in established skills

Age-related dementia is a common concern for most families and caregivers of adults with Down syndrome due to its impact on quality of life. Dementia is more common after age 40 in Down syndrome.^{32,33} However, not everyone gets dementia as they get older. There is no way to diagnose dementia for sure. Some conditions and symptoms commonly mistaken for dementia include cerebrovascular (for example: stroke, aneurysms, narrowing of blood vessels, and moyamoya disease) disease, severe sleep apnea, metabolism problems or medication side effects.^{31, 34}

DEMENTIA IS VERY RARE BEFORE AGE 40 IN PEOPLE WITH DOWN SYNDROME AND CHANGES BEFORE THAT AGE ARE MORE LIKELY TO BE DUE TO ANOTHER CONDITION.

We believe that an accurate diagnosis of dementia is based on more than just changes in behavior or personality. There will usually be changes in more than one area, including memory, self-control, behavior, personality, language, communication, daily living activities, continence (bladder control), and sleep patterns. Other signs of dementia in adults with Down syndrome include seizures, twitching, uncontrolled muscle movements, or loss of walking ability.

3

DIABETES



RECOMMENDATION 5

For asymptomatic adults with Down syndrome, screening for type 2 diabetes mellitus (T2DM) using hemoglobin A1c (HbA1c) or fasting plasma glucose should be performed every 3 years beginning at age 30.

RECOMMENDATION 6

For any adult with Down syndrome and comorbid obesity, screening for T2DM using HbA1c or fasting plasma glucose should be performed every 2-3 years beginning at age 21.


Type 2 diabetes occurs when a person's body has difficulty using sugar (glucose). In order to regulate their sugar levels, they need to take medication and be very careful with their diet. Diabetes is a common concern for adults with Down syndrome, their families, and caregivers since many people with Down syndrome are obese, which can cause diabetes. One study found that adults with Down syndrome are much more likely to have diabetes than adults without Down syndrome.³⁵ We believe that the benefits of screening adults with Down syndrome for diabetes to avoid other complications, such as brain, eye or kidney problems, are greater than the potential harms from diabetes treatment (such as medication side effects).

The American Diabetes Association (ADA) recommends screening all adults for diabetes beginning at age 45.³⁶ However, since adults with Down syndrome tend to age faster and get some conditions (e.g., cataracts) earlier, we recommend that all adults with Down syndrome get screening every 3 years beginning at age 30 for diabetes.³⁷⁻³⁹ Also, obese adults with Down syndrome should be screened even younger and more frequently, every 2-3 years beginning at age 21. Earlier screening may result in more blood draws and over-treatment of diabetes after diagnosis. However, we believe the benefits are greater than the consequences of undiagnosed diabetes.

4

CARDIOVASCULAR DISEASE





RECOMMENDATION 7

For adults with Down syndrome without a history of atherosclerotic cardiovascular disease (ASCVD), the appropriateness of statin therapy should be assessed every 5 years starting at age 40 and using a 10-year risk calculator as recommended for adults without Down syndrome by the U.S. Preventive Services Task Force (USPSTF).⁴⁰

Heart disease is the leading cause of death in the United States, and atherosclerotic cardiovascular disease (ASCVD) is the top cause of heart disease.⁴¹ ASCVD is a build-up of cholesterol or fat in the arteries, which makes it hard for blood to flow. There isn't much data regarding ASCVD in people with Down syndrome. However, the available data shows adults with Down syndrome have a lower risk of heart disease, which is consistent with our experience.^{35, 42} The USPSTF recommends that all adults aged 40-75 get screened for hyperlipidemia (fat in the blood including cholesterol) to assess the risk of heart disease.⁴⁰ Even though adults with Down syndrome are less likely to get heart disease, there isn't enough research to say that adults with Down syndrome should be treated differently than adults without Down syndrome in this regard. Your provider can calculate the risk with their patient by using a 10-year cardiac risk calculator. Patients can also calculate the risk directly since these calculators are available on the internet and as apps. Calculating the risk can help guide decisions such as whether to take a medication to lower cholesterol. The USPSTF recommendation for measuring cholesterol and determining risk ends at age 75 because we do not know for sure if it is helpful to treat people after age 75 or people with a limited life expectancy.⁴³ Since adults with Down syndrome have a shorter life expectancy (60 years) than adults without Down syndrome, stopping screening at a younger age than adults without Down syndrome should be considered. Other risk factors for ASCVD in people with Down syndrome are not yet clear. Unlike adults without Down syndrome, there is no evidence that family history of ASCVD is a risk factor of adults with Down syndrome. In addition, while obesity is more common, ASCVD is less common in people with Down syndrome, suggesting other factors may be more significant in ASCVD in people with Down syndrome.



RECOMMENDATION 8

For adults with Down syndrome, risk factors for stroke should be managed as specified by the American Heart Association/American Stroke Association's Guidelines for the Primary Prevention of Stroke.⁴⁴

RECOMMENDATION 9

In adults with Down syndrome with a history of congenital heart disease, given the elevated risk of cardioembolic stroke, a periodic cardiac evaluation and a corresponding monitoring plan should be reviewed by a cardiologist.

Congenital heart disease is more common in people with Down syndrome. About 50% of children with Down syndrome are born with congenital heart disease.^{45,46} This may include “holes in the heart” or abnormalities of the heart valves. Adults with Down syndrome who had congenital heart disease (whether it was repaired or not) are at a higher risk for a stroke.⁴² A stroke causes damage to the brain due to the blockage of blood flow or the bursting of a blood vessel in the brain. One study found the average age for the first stroke was 41.8 years in those with Down syndrome and 57.1 years in those without Down syndrome.⁴²

Although congenital heart disease is more common, some other risk factors for stroke are less common including hypertension (high blood pressure) and atherosclerosis.⁴⁷⁻⁴⁹ Since the data are limited regarding the risk of stroke for adults with Down syndrome without a history of congenital heart disease, we recommend screening for stroke risk for adults with Down syndrome similarly to those without Down syndrome. However, since those with Down syndrome with congenital heart disease have a higher risk, we recommend additional screening and precaution via appropriate assessment, monitoring, and treatment of the risk factors, even though there is not yet research completed on whether this will reduce strokes. Each person being evaluated and treated for risk factors and strokes is different. Individual risk factors should be discussed with the adult with Down syndrome, their family, caregivers, and health care provider.

5

OBESEITY



★ **Statement of Good Practice 3** Healthy diet, regular exercise, and calorie management should be followed by all adults with Down syndrome as part of a comprehensive approach to weight management, appetite control, and enhancement of quality of life.

Obesity is generally considered having a body mass index (BMI) equal to or greater than 30. BMI is a measure of body proportion based on a person's weight and height. Of course, since people with Down syndrome tend to be shorter and may gain weight more easily, they may have a higher chance of their BMI being greater than 30. Many adults with Down syndrome are obese, so the health consequences of obesity are a common concern for adults with Down syndrome, their families, and caregivers. In order to have good health and quality of life, it is very important to have a healthy diet, portion control, have meals at the same time every day and get regular exercise. We also recommend getting daily exercise that includes socializing with others. Adults with Down syndrome, their families, caretakers, and medical professionals should make decisions together about planning their food and exercise choices.

RECOMMENDATION 10

Monitoring for weight change and obesity should be performed annually by calculating body mass index (BMI) in adults with Down syndrome. The U.S. Preventive Services Task Force (USPSTF) *Behavioral Weight Loss Interventions to Prevent Obesity-Related Morbidity and Mortality in Adults* should be followed.⁵⁰


Many adults with Down syndrome are obese, so the health consequences of obesity are a common concern for adults with Down syndrome, their families, and caregivers. We did not find any specific studies on the effects of obese adults with Down syndrome following a healthy diet, practicing portion control and exercising regularly. This is because most studies focus on exercise only and not on portion control, or focus on both for adults without Down syndrome.⁵¹ The studies that focused on exercise only for adults with Down syndrome showed that it improved their balance, muscle strength and endurance.⁵²

Some factors that contribute to obesity in adults with Down syndrome are lack of physical activity and nutritional guidance, and lack of control over their environment and food choices. Some medical issues and medications also contribute to gaining weight.^{53,54} Adults with Down syndrome face many obstacles when it comes to getting enough exercise,⁵⁵ but not all of them become obese. Families can help their family member with Down syndrome keep healthy and/or lose weight. We have noticed that social group activities like dancing, Zumba, working with a personal trainer, or team sports through the Special Olympics have been very successful.

6

ATLANTOAXIAL INSTABILITY





RECOMMENDATION 11

In adults with Down syndrome, routine cervical spine X-rays should not be used to screen for risk of spinal cord injury in asymptomatic individuals. Instead, annual screening of adults with Down syndrome should include a review of signs and symptoms of cervical myelopathy using targeted history and physical exam.

Atlantoaxial instability (AAI) occurs when the bones at the base of the skull and top of the neck are unstable. About 10% of adults with Down syndrome are found to have AAI when they are X-rayed, but they don't seem to have spinal cord injuries. There are no studies that prove there is a connection between having AAI as seen on an X-ray and having spinal cord injuries. Looking for signs of spinal cord injury during a physical exam is low cost and low risk. There are not many cases of spinal cord injury in people with Down syndrome due to AAI. Special Olympics organizers have reported no spinal cord injuries from over 50,000 individuals with Down syndrome participating in Special Olympics activities over 20 years.⁵⁶

It is important to avoid a spinal cord injury because it can cause other medical problems and even death. However, X-ray findings are not specific and could result in restricting activity unnecessarily. A physical exam should be the first step if they don't show symptoms of spinal cord injury. Some adults with Down syndrome and families/caregivers may feel differently, so it's important to make the decision together. If there are symptoms or signs of spinal cord injury, then X-rays and other tests should be run. Some symptoms of spinal cord injury include walking differently, weaker legs or hands (dropping objects), incontinence, and difficulty holding ones head up.

7

OSTEOPOROSIS



RECOMMENDATION 12

For primary prevention of osteoporotic fractures in adults with Down syndrome, there is insufficient evidence to recommend for or against applying established osteoporosis screening guidelines, including fracture risk estimation; thus, good clinical practice would support a shared decision-making approach to this issue.

Osteoporosis is a bone disease that gets more severe as a person gets older. It happens when the bones lose minerals and become weaker, which causes the bones to break more easily. There is not enough evidence on how to prevent osteoporosis in adults with Down syndrome, and the data available is from adults without Down syndrome. The most common tools for predicting bone breakage (DEXA Scan) were designed for adults without Down syndrome, so they may not work well for adults with Down syndrome. Additionally, we don't know much about how or how many adults with Down syndrome break their bones due to fragility, but it may be very different from adults without Down syndrome.

The most common drugs for preventing and treating osteoporosis may not be the most effective, and may even be dangerous for adults with Down syndrome.⁵⁷ We are concerned that diagnosing and treating adults with Down syndrome for osteoporosis the same way as adults without Down syndrome could cause problems, so we recommend that each family makes the decision together with their medical professional on the potential risks, benefits and uncertainties around osteoporosis screening and prevention, including medications, exercise, and vitamin D.

WE DO RECOMMEND MUSCLE STRENGTHENING EXERCISES, AS THEY ARE GOOD FOR BONE HEALTH AND ARE USUALLY SAFE FOR ADULTS WITH DOWN SYNDROME.⁵²





RECOMMENDATION 13

All adults with Down syndrome who sustain a fragility fracture should be evaluated for secondary causes of osteoporosis, including screening for hyperthyroidism, celiac disease, vitamin D deficiency, hyperparathyroidism and medications associated with adverse effects on bone health.

Potential secondary causes of osteoporosis are more prevalent in adults with Down syndrome than in adults without Down syndrome. Lack of exercise and certain medical conditions can lead to adults with Down syndrome breaking a bone due to osteoporosis.⁵⁸⁻⁶² Many of these conditions can be corrected and bone health can be improved with healthy habits. Researchers are studying if vitamin D supplementation works for preventing and treating osteoporosis in adults without Down syndrome.⁶³ It is common for adults with Down syndrome to have low vitamin D. Vitamin D might help their bones and doesn't seem to cause harm.⁶⁴ In addition to vitamin D, we recommend looking at all options for preventing osteoporosis as is recommended for adults without Down syndrome, including getting enough calcium, preventing falls, improving vision, and lifting weights.⁶⁵

8

THYROID



RECOMMENDATION 14

Screening adults with Down syndrome for hypothyroidism should be performed every 1-2 years using a serum thyroid-stimulating hormone (TSH) test beginning at age 21.

The thyroid gland is located in the neck and produces hormones related to activity, metabolism and heart rate. If the thyroid makes too much hormone, that is called hyperthyroidism. If the thyroid makes too little hormone, that is called hypothyroidism. Three studies show that adults with Down syndrome are more likely to have thyroid problems than adults without Down syndrome.^{35, 66, 67} It is common for adults with Down syndrome to have hypothyroidism and is more common as they get older.

DIAGNOSING HYPOTHYROIDISM IN ADULTS WITH DOWN SYNDROME CAN BE HARD BECAUSE SOME OF THE SYMPTOMS, FOR EXAMPLE FATIGUE, WEIGHT GAIN AND CONSTIPATION, ARE COMMON IN ADULTS WITH DOWN SYNDROME WHETHER OR NOT THEY HAVE THYROID PROBLEMS.

Relying only on symptoms is not enough, so we believe it is important to use a TSH blood test every 1-2 years. Even though having a blood draw can cause anxiety, getting a correct diagnosis could prevent future behavior changes and weight gain, which are hard to reverse later. Proper treatment can also improve constipation, dry skin, fatigue and more. Treatment is inexpensive, safe, and works better if hypothyroidism is detected early instead of waiting for obvious signs.

9

CELIAC DISEASE





★ **Statement of Good Practice 4** Adults with Down syndrome should receive an annual assessment for gastrointestinal and non-gastrointestinal signs and symptoms of celiac disease using targeted history, physical examination, and clinical judgment of good practice.

Celiac disease is an autoimmune condition where your body has a reaction to gluten (a protein found in foods containing wheat, rye and barley). Studies show that celiac disease is more common in children with Down syndrome.^{68, 69} Adult onset of celiac disease, however, has not been well studied in adults with Down syndrome. Diagnosing celiac disease in adults with Down syndrome may be challenging due to communication difficulties and the fact that signs and symptoms of celiac disease can be common symptoms of other conditions. Also, many of the signs and symptoms, like constipation or rashes, are common in Down syndrome, even if they don't have celiac disease. Therefore, in addition to stomach problems, medical professionals should evaluate for celiac disease every year using a physical exam paying attention to behavioral changes, rashes, and symptoms of other possible autoimmune disorders. Confirming a diagnosis of celiac disease requires laboratory testing and a bowel biopsy, so some families who suspect celiac disease may try a gluten-free diet at home to see if it helps.

GLOBAL DOWN SYNDROME FOUNDATION & AFFILIATES

GLOBAL DOWN SYNDROME FOUNDATION. The Global Down Syndrome Foundation (GLOBAL) is the largest non-profit in the U.S. working to save lives and dramatically improve health outcomes for people with Down syndrome. GLOBAL established the first Down syndrome research institute supporting over 400 scientists and over 2,200 patients with Down syndrome from 33 states and 10 countries. Working closely with Congress and the National Institutes of Health, GLOBAL is the lead advocacy organization in the U.S. for Down syndrome research and care. GLOBAL has a membership of over 120 Down syndrome organizations worldwide and is part of a network of Affiliates.

FOR MORE INFORMATION VISIT OUR WEBSITE: <https://www.globaldownsyndrome.org/>

GLOBAL DOWN SYNDROME FOUNDATION MEDICAL CARE GUIDELINES FOR ADULTS WITH DOWN SYNDROME

<https://www.globaldownsyndrome.org/medical-care-guidelines-for-adults/>

DOWN SYNDROME MEDICAL CARE CENTERS IN THE U.S. BY STATE

<https://www.globaldownsyndrome.org/research-medical-care/medical-care-providers/>

LOCAL DOWN SYNDROME ORGANIZATIONS IN THE U.S. BY STATE

<https://www.globaldownsyndrome.org/about-down-syndrome/resources/local-organizations/>

LINDA CRNIC INSTITUTE FOR DOWN SYNDROME THE WORLD'S LEADING DOWN SYNDROME RESEARCH INSTITUTE

<https://medschool.cuanschutz.edu/linda-crnic-institute>

ANNA AND JOHN J. SIE CENTER FOR DOWN SYNDROME ONE OF THE WORLD'S LARGEST DOWN SYNDROME MEDICAL CARE CENTERS

<https://www.globaldownsyndrome.org/our-story/anna-and-john-j-sie-center-for-down-syndrome>

UNIVERSITY OF COLORADO ALZHEIMER'S AND COGNITION CENTER

<https://medschool.cuanschutz.edu/alzheimer>

** In loving memory of Dr. Kent McKelvey, who dedicated his life to caring for adults with Down syndrome. His leadership and belief in GLOBAL gave us the confidence to create this important guideline. We are forever grateful.*



ADDITIONAL DOWN SYNDROME RESOURCES

AMERICAN ACADEMY OF DEVELOPMENTAL MEDICINE AND DENTISTRY

<https://www.aadmd.org/>

DOWN SYNDROME-AUTISM CONNECTION

<http://www.ds-asd-connection.org/>

DOWN SYNDROME MEDICAL INTEREST GROUP-USA

<https://www.dsmig-usa.org/>

DS CONNECT® AN IMPORTANT REGISTRY AND PORTAL CONNECTING PEOPLE WITH DOWN SYNDROME, CLINICIANS, AND RESEARCHERS

<https://dsconnect.nih.gov/>

INCLUDE DATA COORDINATING CENTER

<https://includedcc.org/>

INTERNATIONAL MOSAIC DOWN SYNDROME ASSOCIATION

<https://www.imdsa.org/mosaic-down-syndrome>

NATIONAL DOWN SYNDROME CONGRESS

<https://www.ndsccenter.org/>

NATIONAL DOWN SYNDROME SOCIETY

<https://www.ndss.org/>

NATIONAL INSTITUTES OF HEALTH DOWN SYNDROME CONSORTIUM

<https://downsyndrome.nih.gov/>

NATIONAL TASK GROUP ON INTELLECTUAL DISABILITIES AND DEMENTIA PRACTICES

<https://www.the-ntg.org/>

SPECIAL OLYMPICS

<https://www.specialolympics.org/>

THE ARC

<https://thearc.org/>

REFERENCES

1. American Academy of Pediatrics Committee on Genetics. Health supervision for children with Down syndrome. *Pediatrics*. May 1994;93(5):855-9.
2. Bull MJ, Trotter T, Santoro SL, Christensen C, Grout RW, The Council On Genetics. Health supervision for children and adolescents with Down syndrome. *Pediatrics*. May 2022;149(5):e 2022057010. doi:10.1542/peds.2022-057010
3. Cohen WL. Health care guidelines for individuals with Down syndrome. *Down Syndrome Quarterly*. 3. 1999;4:1-16.
4. Smith DS. Health care management of adults with Down syndrome. *Am Fam Physician*. Sep 2001;64(6):1031-8.
5. Tsou AY, Bulova P, Capone G, et al. Medical care of adults with Down syndrome: A clinical guideline. *JAMA*. 2020;324(15):1543-1556. doi:10.1001/jama.2020.17024
6. Shin M, Siffel C, Correa A. Survival of children with mosaic Down syndrome. *Am J Med Genet A*. Mar 2010;152A(3):800-1. doi:10.1002/ajmg.a.33295
7. Mai CT, Isenburg JL, Canfield MA, et al. National population-based estimates for major birth defects, 2010-2014. *Birth Defects Res*. 11 2019;111(18):1420-1435. doi:10.1002/bdr2.1589
8. Parker SE, Mai CT, Canfield MA, et al. Updated National Birth Prevalence estimates for selected birth defects in the United States, 2004-2006. *Birth Defects Res A Clin Mol Teratol*. Dec 2010;88(12):1008-16. doi:10.1002/bdra.20735
9. National Human Genome Research Institute. About Down syndrome. <https://www.genome.gov/Genetic-Disorders/Down-Syndrome> Accessed August 27, 2019.
10. Hasle H, Friedman JM, Olsen JH, Rasmussen SA. Low risk of solid tumors in persons with Down syndrome. *Genet Med*. 11 2016;18(11):1151-1157. doi:10.1038/gim.2016.23
11. Parra P, Costa R, de Asúa DR, Moldenhauer F, Suárez C. Atherosclerotic surrogate markers in adults with Down syndrome: A case-control study. *J Clin Hypertens (Greenwich)*. Feb 2017;19(2):205-211. doi:10.1111/jch.12890
12. Hickey F, Hickey E, Summar KL. Medical update for children with Down syndrome for the pediatrician and family practitioner. *Adv Pediatr*. 2012;59(1):137-57. doi:10.1016/j.yapd.2012.04.006
13. Weijerman M, JP dW. Clinical practice. The care of children with Down syndrome. *European Journal of Pediatrics*. 2010;169(12):1445-1452. doi:10.1007/s00431-010-1253-0.
14. Bull MJ. Down Syndrome. *N Engl J Med*. 06 2020;382(24):2344-2352. doi:10.1056/NEJMra1706537
15. Shin M, Besser LM, Kucik JE, et al. Prevalence of Down syndrome among children and adolescents in 10 regions of the United States. *Pediatrics*. Dec 2009;124(6):1565-71. doi:10.1542/peds.2009-0745
16. de Graaf G, Buckley F, Skotko BG. Estimation of the number of people with Down syndrome in the United States. *Genetics in Medicine*. Sep 08 2017:439-447. doi:10.1038/gim.2016.127
17. Steingass KJ, Chicoine B, McGuire D, Roizen NJ. Developmental disabilities grown up: Down syndrome. *Journal of Developmental and Behavioral Pediatrics*. Sep 2011;32(7):548-558. doi:10.1097/DBP.0b013e31822182e0
18. Pai M, Santesso N, Yeung CH, Lane SJ, Schünemann HJ, Iorio A. Methodology for the development of the NHF-McMaster Guideline on Care Models for Haemophilia Management. *Haemophilia*. Jul 2016;22 Suppl 3:17-22. doi:10.1111/hae.13007
19. Vicari S, Pontillo M, Armando M. Neurodevelopmental and psychiatric issues in Down's syndrome: Assessment and intervention. *Psychiatr Genet*. Jun 2013;23(3):95-107. doi:10.1097/YPG.0b013e32835fe426

REFERENCES

20. Capone G, Aidikoff J, Taylor K, Rykiel N. Adolescents and young adults with down syndrome presenting to a medical clinic with depression: Co-morbid obstructive sleep apnea. *American Journal of Medical Genetics, Part A*. Sep 2013;161(9):2188-2196.
21. Jacobs J, Schwartz A, McDougle CJ, Skotko BG. Rapid clinical deterioration in an individual with Down syndrome. *Am J Med Genet A*. 07 2016;170(7):1899-902. doi:10.1002/ajmg.a.37674
22. Mircher C, Cieuta-Walti C, Marey I, et al. Acute regression in young people with Down syndrome. *Brain Sci*. Jun 2017;7(6)doi:ARTN 5710.3390/brainsci7060057
23. Chicoine B, Capone G. Regression in Adolescents and Adults with Down Syndrome. In: Prasher VP, Janicki MP, eds. *Physical Health of Adults with Intellectual and Developmental Disabilities*. Springer, Cham; 2018:121-140:chap Regression in Adolescents and Adults with Down Syndrome.
24. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders V In*. Washington, D.C.: American Psychiatric Association; 2013
25. Fletcher R. *Diagnostic manual- intellectual disability: A clinical guide for diagnosis of mental disorders in persons with intellectual disability*. 2nd edition. ed. NADD Press; 2018:pages cm.
26. Glenn S, Cunningham C, Nananidou A, Prasher V, Glenholmes P. Routinised and compulsive-like behaviours in individuals with Down syndrome. *J Intellect Disabil Res*. Nov 2015;59(11):1061-70. doi:10.1111/jir.12199
27. Foley KR, Bourke J, Einfeld SL, Tonge BJ, Jacoby P, Leonard H. Patterns of depressive symptoms and social relating behaviors differ over time from other behavioral domains for young people with Down syndrome. *Medicine (Baltimore)*. May 2015;94(19):e710. doi:10.1097/md.0000000000000710
28. Straccia C, Baggio S, Barisnikov K. Mental illness, behavior problems, and social behavior in adults with Down syndrome. research-article. <http://dxdoiorg/101080/193158642012741660>. 3 Dec 2013 2013;doi:Journal of Mental Health Research in Intellectual Disabilities, Vol. 7, No. 1, January-March 2014: pp. 74–90
29. Fletcher R, Havercamp S, Ruedrich S, et al. Clinical usefulness of the diagnostic manual-intellectual disability for mental disorders in persons with intellectual disability: results from a brief field survey. *J Clin Psychiatry*. Jul 2009;70(7):967-74. doi:10.4088/JCP.08m04429
30. Fletcher RJ, National Association for the Dually D, American Psychiatric A. *DM-ID: diagnostic manual-intellectual disability: a clinical guide for diagnosis of mental disorders in persons with intellectual disability*. NADD Press; 2007.
31. Moran JA, Rafi MS, Keller SM, Singh BK, Janicki MP. The National Task Group on Intellectual Disabilities and Dementia Practices consensus recommendations for the evaluation and management of dementia in adults with intellectual disabilities. *Mayo Clin Proc*. Aug 2013;88(8):831-40. doi:10.1016/j.mayocp.2013.04.024
32. Coppus A, Evenhuis H, Verberne GJ, et al. Dementia and mortality in persons with Down's syndrome. *J Intellect Disabil Res*. Oct 2006;50(Pt 10):768-77. doi:10.1111/j.1365-2788.2006.00842.x
33. Fortea J, Vilaplana E, Carmona-Iragui M, et al. Clinical and biomarker changes of Alzheimer's disease in adults with Down syndrome: A cross-sectional study. *Lancet*. 06 2020;395(10242):1988-1997. doi:10.1016/S0140-6736(20)30689-9
34. Prasher VP, Mahmood H, Mitra M. Challenges faced in managing dementia in Alzheimer's disease in patients with Down syndrome. *Degener Neurol Neuromuscul Dis*. 2016;6:85-94. doi:10.2147/dnnd.s91754

REFERENCES

35. Alexander M, Petri H, Ding Y, Wandel C, Khwaja O, Foskett N. Morbidity and medication in a large population of individuals with Down syndrome compared to the general population. *Dev Med Child Neurol*. Mar 2016;58(3):246-54. doi:10.1111/dmcn.12868
36. American Diabetes Association. 2. Classification and Diagnosis of Diabetes. *Diabetes Care*. Jan 2020;43(Suppl1):S14-S31. doi:10.2337/dc20-S002
37. Krinsky-McHale SJ, Jenkins EC, Zigman WB, Silverman W. Ophthalmic disorders in adults with Down syndrome. *Curr Gerontol Geriatr Res*. 2012;2012:974253. doi:10.1155/2012/974253
38. Patel A, Yamashita N, Ascaño M, et al. RCAN1 links impaired neurotrophin trafficking to aberrant development of the sympathetic nervous system in Down syndrome. *Nat Commun*. Dec 2015;6:10119. doi:10.1038/ncomms10119
39. Lo A, Brown HG, Fivush BA, Neu AM, Racusen LC. Renal disease in Down syndrome: Autopsy study with emphasis on glomerular lesions. *Am J Kidney Dis*. Feb 1998;31(2):329-35. doi:10.1053/ajkd.1998.v31.pm9469506
40. Bibbins-Domingo K, Grossman DC, Curry SJ, et al. Statin use for the primary prevention of cardiovascular disease in adults: US Preventive Services Task Force Recommendation Statement. *JAMA*. Nov 15 2016;316(19):1997-2007. doi:10.1001/jama.2016.15450
41. Centers for Disease Control and Prevention. Heart Disease Facts. Updated June 22, 2020. <https://www.cdc.gov/heartdisease/facts.htm>. Accessed February 12, 2019.
42. Sobey CG, Judkins CP, Sundararajan V, Phan TG, Drummond GR, Srikanth VK. Risk of major cardiovascular events in people with Down syndrome. *PLoS One*. 2015;10(9):e0137093. doi:10.1371/journal.pone.0137093
43. The Society for Post-Acute and Long-Term Care Medicine. Cholesterol drugs for people 75 and older: When you need them -- and when you don't. Updated 2017. <https://www.choosingwisely.org/wp-content/uploads/2018/02/Cholesterol-Drugs-For-People-75-And-Older-AMDA.pdf>. Accessed August 11, 2020.
44. Meschia JF, Bushnell C, Boden-Albala B, et al. Guidelines for the primary prevention of stroke: A statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. Dec 2014;45(12):3754-832. doi:10.1161/str.0000000000000046
45. Lanz J, Brophy JM, Therrien J, Kaouache M, Guo L, Marelli AJ. Stroke in adults with congenital heart Disease: Incidence, cumulative risk, and predictors. *Circulation*. Dec 2015;132(25):2385-94. doi:10.1161/CIRCULATIONAHA.115.011241
46. Centers for Disease Control and Prevention. Facts About Down Syndrome. <https://www.cdc.gov/ncbddd/birthdefects/downsyndrome.html>. Accessed April 2019.
47. Capone G, Chicoine B, Bulova P, et al. Co-occurring medical conditions in adults with Down syndrome: A systematic review toward the development of health care guidelines. *Am J Med Genet A*. Jan 2018;176(1):116-133. doi:10.1002/ajmg.a.38512
48. Kainth DS, Chaudhry SA, Kainth HS, Suri FK, Qureshi AI. Prevalence and characteristics of concurrent Down syndrome in patients with moyamoya disease. *Neurosurgery*. Feb 2013;72(2):210-5; discussion 215. doi:10.1227/NEU.0b013e31827b9beb
49. Maris M, Verhulst S, Wojciechowski M, Van de Heyning P, Boudewyns A. Prevalence of obstructive sleep apnea in children with Down syndrome. *Sleep*. Mar 1 2016;39(3):699-704. doi:10.5665/sleep.5554
50. Curry SJ, Krist AH, Owens DK, et al. Behavioral weight loss interventions to prevent obesity-related morbidity and mortality in adults: US Preventive Services Task Force Recommendation Statement. *JAMA*. 09 2018;320(11):1163-1171. doi:10.1001/jama.2018.13022
51. Clark JE. Diet, exercise or diet with exercise: Comparing the effectiveness of treatment options for weight-loss and changes in fitness for adults (18-65 years old) who are overfat, or obese; systematic review and meta-analysis. *J Diabetes Metab Disord*. 2015;14:31. doi:10.1186/s40200-015-0154-1


REFERENCES

52. Li C, Chen S, Meng How Y, Zhang AL. Benefits of physical exercise intervention on fitness of individuals with Down syndrome: a systematic review of randomized-controlled trials. *Int J Rehabil Res*. Sep 2013;36(3):187-95. doi:10.1097/MRR.0b013e3283634e9c
53. Allison DB, Mentore JL, Heo M, et al. Antipsychotic-induced weight gain: a comprehensive research synthesis. *Am J Psychiatry*. Nov 1999;156(11):1686-96. doi:10.1176/ajp.156.11.1686
54. Bertapelli F, Pitetti K, Agiovlasitis S, Guerra-Junior G. Overweight and obesity in children and adolescents with Down syndrome-prevalence, determinants, consequences, and interventions: A literature review. *Research in Developmental Disabilities*. Oct 2016;57:181-192. doi:10.1016/j.ridd.2016.06.018
55. Mendonca GV, Pereira FD, Fernhall B. Cardiac autonomic function during submaximal treadmill exercise in adults with Down syndrome. *Res Dev Disabil*. Mar-Apr 2011;32(2):532-9. doi:S0891-4222(10)00318-5 [pii] 10.1016/j.ridd.2010.12.028
56. Wisconsin Special Olympics. Special Olympics official policy affecting athletes with Down syndrome. 2015. <http://www.specialolympicswisconsin.org/wp-content/uploads/2015/04/Athletes-with-Down-Syndrome-Special-Examination-Form.pdf>. Accessed August 19, 2020.
57. LaCombe JM, Roper RJ. Skeletal dynamics of Down syndrome: A developing perspective. *Bone*. Apr 2020;133:115215. doi:10.1016/j.bone.2019.115215
58. Heller T, Hsieh K, Rimmer J. Barriers and supports for exercise participation among adults with Down syndrome. *Journal of Gerontological Social Work*. 2002;38:161-178.
59. Attia A, Ghanayem N, El Naqeeb H. Sexual and reproductive functions in men with Down's syndrome. *Menoufia Medical Journal*. 2015;28(2):471-476.
60. Schupf N, Zigman W, Kapell D, Lee JH, Kline J, Levin B. Early menopause in women with Down's syndrome. *J Intellect Disabil Res*. Jun 1997;41 (Pt 3):264-7. doi:10.1111/j.1365-2788.1997.tb00706.x
61. Rosello L, Torres R, Boronat T, Llobet R, Puerto E. Osteoporosis prevalence in a Down syndrome population, measuring different parameters. *SD Revista Medica Internacional sobre el Sindrome de Down*; 2004. p. 18-22.
62. Du Y, Shan LF, Cao ZZ, Feng JC, Cheng Y. Prevalence of celiac disease in patients with Down syndrome: A meta-analysis. *Oncotarget*. Jan 2018;9(4):5387-5396. doi:10.18632/oncotarget.23624
63. Lips P, van Schoor NM. The effect of vitamin D on bone and osteoporosis. *Best Pract Res Clin Endocrinol Metab*. Aug 2011;25(4):585-91. doi:10.1016/j.beem.2011.05.002
64. Zubillaga P, Garrido A, Mugica I, Ansa J, Zabalza R, Emparanza JI. Effect of vitamin D and calcium supplementation on bone turnover in institutionalized adults with Down's Syndrome. *Eur J Clin Nutr*. May 2006;60(5):605-9. doi:10.1038/sj.ejcn.1602357
65. Fares A. Pharmacological and non-pharmacological means for prevention of fractures among elderly. *Int J Prev Med*. 2018;9:78. doi:10.4103/ijpvm.IJPVM_114_18
66. Real de Asua D, Quero M, Moldenhauer F, Suarez C. Clinical profile and main comorbidities of Spanish adults with Down syndrome. *European Journal of Internal Medicine*. Jul 2015;26(6):385-391. doi:10.1016/j.ejim.2015.05.003
67. Kerins G, Petrovic K, Bruder MB, Gruman C. Medical conditions and medication use in adults with Down syndrome: A descriptive analysis. *Downs Syndr Res Pract*. Oct 2008;12(2):141-7. doi:10.3104/reports.2009
68. Csizmadia CG, Mearin ML, Oren A, et al. Accuracy and cost-effectiveness of a new strategy to screen for celiac disease in children with Down syndrome. *J Pediatr*. Dec 2000;137(6):756-61. doi:10.1067/mpd.2000.110421
69. Sharr C, Lavigne J, Elsharkawi IM, et al. Detecting celiac disease in patients with Down syndrome. *Am J Med Genet A*. 12 2016;170(12):3098-3105. doi:10.1002/ajmg.a.37879


Image on page 36 courtesy of Logan Rose Photography, Inc.

All images are approved GLOBAL models.

ADDITIONAL GLOBAL RESOURCES




GLOBAL Adult Care Guideline for Clinicians



GLOBAL Family-Friendly Adult Care Guideline



GLOBAL Adult Care Guideline Checklist



COVID-19 & Down Syndrome Resource



Prenatal Testing & Info About Down Syndrome



Award-Winning Down Syndrome World Magazine



OUR COMMITMENT

GLOBAL is committed to working with our Down syndrome community – our self-advocates, our families, our members, and our medical experts – to ensure that the GLOBAL Adult Guideline is updated and published every six years.

We can only publish this important resource with the support from our generous donors and knowing that our families will use the guidelines to advocate for the best possible health outcomes.

Together we are creating a brighter future for our loved ones with Down syndrome!





GLOBAL DOWN SYNDROME FOUNDATION®

The Global Down Syndrome Foundation (GLOBAL) is the largest non-profit in the U.S. working to save lives and dramatically improve health outcomes for people with Down syndrome. GLOBAL has donated more than \$32 million to establish the first Down syndrome research institute supporting over 400 scientists and over 2,200 patients with Down syndrome from 33 states and 10 countries. Working closely with Congress and the National Institutes of Health, GLOBAL is the lead advocacy organization in the U.S. for Down syndrome research and medical care. GLOBAL has a membership of over 120 Down syndrome organizations worldwide and is part of a network of Affiliates – the Crnic Institute for Down Syndrome, the Sie Center for Down Syndrome, and the University of Colorado Alzheimer's and Cognition Center – all on the Anschutz Medical Campus. GLOBAL's widely circulated medical publications include *GLOBAL Medical Care Guidelines for Adults with Down Syndrome*, *Prenatal Testing and Information about Down Syndrome*, and the award-winning *Down Syndrome World™* magazine. GLOBAL also organizes the *Be Beautiful Be Yourself Fashion Show*, the largest Down syndrome fundraiser in the world.

GLOBAL DOWN SYNDROME FOUNDATION

3239 E 2ND AVE
DENVER, CO 80206-5203

GLOBALDOWNSYNDROME.ORG